

CONTENTS

VOLLETH, M., J. LOIDL, F. MAYER, H.-S. YONG, S. MÜLLER, and K.-G. HELLER Surprising genetic diversity in <i>Rhinolophus luctus</i> (Chiroptera: Rhinolophidae) from Peninsular Malaysia: description of a new species based on genetic and morphological characters	1
SOISOOK, P., M. J. STRUEBIG, S. NOERFAHMY, H. BERNARD, I. MARYANTO, S.-F. CHEN, S. J. ROSSITER, H.-C. KUO, K. DESHPANDE, P. J. J. BATES, D. SYKES, and R. PORTELA MIGUEZ Description of a new species of the <i>Rhinolophus trifolius</i> -group (Chiroptera: Rhinolophidae) from Southeast Asia	21
NÁJERA-CORTAZAR, L. A., S. T. ÁLVAREZ-CASTAÑEDA, and E. DE LUNA An analysis of <i>Myotis peninsularis</i> (Vespertilionidae) blending morphometric and genetic datasets	37
KRUSKOP, S. V. Dark and pale: taxonomic status of the barbastelle (<i>Barbastella</i> : Vespertilionidae, Chiroptera) from Central Asia	49
BOLZAN, D. P., L. M. PESSÔA, A. L. PERACCHI, and R. E. STRAUSS Allometric patterns and evolution in Neotropical nectar-feeding bats (Chiroptera, Phyllostomidae)	59
HURTADO, N., R. D. SEPÚLVEDA, and V. PACHECO Sexual size dimorphism of a sensory structure in a monomorphic bat	75
RAMASINDRAZANA, B., C. F. RAKOTONDRAMANANA, M. C. SCHOEMAN, and S. M. GOODMAN Evidence of echolocation call divergence in <i>Hipposideros commersoni</i> sensu stricto (E. Geoffroy, 1803) from Madagascar and correlation with body size	85
RALISATA, M., D. RAKOTONDRAVONY, and P. A. RACEY The relationship between male sucker-footed bats <i>Myzopoda aurita</i> and the traveller's tree <i>Ravenala madagascariensis</i> in south-eastern Madagascar	95
NAKAMOTO, A., K. KINJO, and M. IZAWA Dietary plasticity in the Ryukyu flying fox on a subtropical island at the northern range limit of <i>Pteropus</i>	105
LÓPEZ-GONZÁLEZ, C., E. P. GÓMEZ-RUIZ, A. LOZANO, and R. LÓPEZ-WILCHIS Activity of insectivorous bats associated with cattle ponds at La Michilía Biosphere Reserve, Durango, Mexico: implications for conservation	117
LUCAS, J. S., S. C. LOEB, and P. G. R. JODICE Roost selection by Rafinesque's big-eared bats (<i>Corynorhinus rafinesquii</i>) in a pristine habitat at three spatial scales	131
GIMÉNEZ, A. L., N. P. GIANNINI, M. I. SCHIAFFINI, and G. M. MARTIN Geographic and potential distribution of a poorly known South American bat, <i>Histiotus macrotus</i> (Chiroptera: Vespertilionidae)	143
HAYES, M. A., K. OZENBERGER, P. M. CRYAN, and M. B. WUNDER Not to put too fine a point on it — does increasing precision of geographic referencing improve species distribution models for a wide-ranging migratory bat?	159
McFARLANE, D. A., G. VAN RENTERGEM, A. RUINA, J. LUNDBERG, and K. CHRISTENSON Estimating colony size of the wrinkle-lipped bat, <i>Chaerephon plicatus</i> (Chiroptera: Molossidae) at Gomantong, Sabah, by quantitative image analysis	171
DRAKE, D., C. S. JENNELLE, J.-N. LIU, S. M. GRODSKY, S. SCHUMACHER, and M. SPONSLER Regional analysis of wind turbine-caused bat mortality	179
URBAN, D. J., D. W. SORENSEN, J. A. MAIER, M. B. FENTON, N. B. SIMMONS, L. N. COOPER, and K. E. SEARS Conjoined twins in a wild bat: a case report	189
GORRESEN, P. M., P. M. CRYAN, D. C. DALTON, S. WOLF, and F. J. BONACCORSO Ultraviolet vision may be widespread in bats	193
BOOK REVIEWS	199